

TALKING WITH YOUR HEALTHCARE PROVIDER

HERE ARE SOME TIPS TO HELP YOU HAVE MORE MEANINGFUL DISCUSSIONS

A visit with your healthcare provider can go by pretty fast—and it's easy to leave without saying everything you had planned to say. Your healthcare provider may begin the conversation by asking you about your treatment history, how you are feeling, and any medications you may be taking. However, preparing a list of your own questions can also help. While this is not a complete list, here are some suggestions.

HERE ARE SOME QUESTIONS YOU CAN ASK ABOUT PERSERIS

- How can PERSERIS™ (risperidone) help me manage my schizophrenia?
- What are the possible side effects of PERSERIS?
- Who should not take PERSERIS?
- If PERSERIS is right for me, how soon can I start treatment?

If you have any other questions or concerns, be sure to discuss them with your healthcare provider.

KEEP THE CONVERSATION GOING WITH YOUR HEALTHCARE PROVIDER

After you begin treatment, it's important to keep the lines of communication open with your healthcare provider and any other members of your Treatment Team. The more they know about your health, the better they can help you.

Here are some things you may want to talk about during your monthly visits:

- Your current treatment goals, or setting new goals
- Any changes in your symptoms, your health, or how you have been feeling since your last visit
- How to reach out to others, such as caregivers or patient support groups
- If you have any schedule conflicts that make it difficult to get to your monthly appointments

If you are experiencing side effects at any time, be sure to speak with your healthcare provider as soon as possible.

INDICATION


PERSERIS™ (risperidone) for extended-release injectable suspension, for subcutaneous use, is a prescription medicine used to treat schizophrenia in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about PERSERIS?

Drugs like PERSERIS that are used to treat schizophrenia can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). PERSERIS is not approved to treat dementia-related psychosis and it has not been studied in elderly people with dementia-related psychosis.

Please see additional Important Safety Information throughout this guide and visit [PERSERIS.com](https://www.perseris.com) to see the full Prescribing Information including **BOXED WARNING**, or talk to your healthcare provider.

once-monthly
 **PERSERIS™**
(risperidone)
for extended-release
injectable suspension
90 mg • 120 mg

NOTES

Use the space below to write down any additional questions or concerns you may have.


IMPORTANT SAFETY INFORMATION (CONT.)

Do not receive PERSERIS if you are allergic to risperidone, paliperidone, or any of its components. Allergic reactions, including swelling of the face, mouth, lips, and tongue, have been reported in some patients treated with risperidone.

PERSERIS may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** that may not go away, even if you stop receiving PERSERIS. Tardive dyskinesia may also start after you stop receiving PERSERIS.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with PERSERIS. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.
- **High level of prolactin in your blood.** PERSERIS may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like PERSERIS may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body. Do not** drive, operate machinery, or do other dangerous activities until you know how PERSERIS affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

Please see additional Important Safety Information throughout this guide and visit [PERSERIS.com](https://www.perseris.com) to see the full Prescribing Information including **BOXED WARNING**, or talk to your healthcare provider.

once-monthly
 **PERSERIS™**
 (risperidone)
 for extended-release
 injectable suspension
90 mg • 120 mg

IMPORTANT SAFETY INFORMATION (CONT.)

The most common side effects of PERSERIS include weight gain; constipation; drowsiness; pain in arms, legs, back, or muscles; restlessness; feeling excessive worry or nervousness; and pain or redness at the injection site. These are not all the possible side effects of PERSERIS. Tell your healthcare provider if you have any side effect that bothers you or does not go away. For more information, ask your healthcare provider or pharmacist.

Do not drink alcohol during treatment with PERSERIS.

Before receiving PERSERIS, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.
- have had a low white blood cell count.
- have or have had seizures or epilepsy.
- are pregnant or plan to become pregnant during treatment with PERSERIS. It is not known if PERSERIS will harm your unborn baby. Use of PERSERIS during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with PERSERIS, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.

- are breastfeeding or plan to breastfeed. If you are receiving PERSERIS and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. PERSERIS and other medicines may affect each other.

To report pregnancy or side effects associated with taking PERSERIS, please call 1-877-782-6966. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For more information about PERSERIS and the full Prescribing Information including BOXED WARNING, visit PERSERIS.com or talk to your healthcare provider.



PERSERIS is a trademark of Indivior UK Limited.

© Indivior PLC | INDIVIOR is a registered trademark of Indivior UK Limited | All rights reserved.

PERSERIS is manufactured for Indivior Inc., North Chesterfield, VA 23235 by AMRI, Burlington, MA 01803.

This material is intended solely for residents of the United States.

P-RAG-US-00094 EXPIRY: May 2021

